European Journal of Pharmacology, 181 (1990) 333–335
Elsevier

Author index

Abdelrahman, A., Nguyen, H. and Pang, C.C.Y.
Reversal of α-adrenoceptor blockade by propranolol in isolated rat pulmonary artery (181) 83
Adell, A., see Artigas, F. (181) 9
Akaji, K.-i., see Katsuki, H. (181) 323
Allgaier, C., Choi, B.K. and Hertting, G.
Forskolin modulates acetylcholine release in the hippocampus independently of adenylate cyclase activation (181) 279
Altar, C.A., Boyar, W.C. and Kim, H.S.
Discriminatory roles for D_1 and D_2 dopamine receptor subtypes in the in vivo control of neostriatal cyclic GMP (181) 17
Amann, R., see Donnerer, J. (181) 147
Aniksztejn, L. and Ben-Ari, Y.
NMDA-independent form of long-term potentiation produced by tetraethylamonium in the hippocampal CA1 region (181) 157
Artigas, F., Martínez, E. and Adell, A.
Non-specific inhibition of imipramine binding argues against an endogenous ligand (181) 9
Azrad, A., see Leiba, H. (181) 71

Baba, A., see Matsuda, T. (181) 295
Ben-Ari, Y., see Aniksztejn, L. (181) 157
Bettaney, J., see Dray, A. (181) 289
Black, C.K., see Schmidt, C.J. (181) 59
Bohus, B., see Korte, S.M. (181) 307
Bohus, B.G.J., see Timmerman, W. (181) 253
Boix, F., see Fernández-Teruel, A. (181) 35
Bouhelal, R. and Mir, A.K.
Role of the adrenal gland in the metabolic and cardiovascular effects of 8-OH-DPAT in the rat (181) 89
Boyar, W.C., see Altar, C.A. (181) 17

Canonico, P.L., see Copani, A. (181) 327
Carey, H.V. and Zafirova, M.
Adrenergic inhibition of neurally evoked secretion in ground squirrel intestine (181) 43
Chagraoui, A., see Vasse, M. (181) 23
Choi, B.K., see Allgaier, C. (181) 279
Cohen, A., Weizman, A., Eldar, M. and Rehavi, M.
Long-term verapamil treatment increases [^{3}H]imipramine binding in human platelets (181) 311
Copani, A., Canonico, P.L. and Nicoletti, F.
β-N-Methylamino-L-Alanine (L-BMAA) is a potent agonist of 'metabolotropic' glutamate receptors (181) 327

Den Hertog, A., see Hoiting, B. (181) 127
DiMicco, J.A., see Wible, Jr., J.H. (181) 215
Donnerer, J., Amann, R., Skofitsch, G. and Lembeck, F.
Region-specific noradrenaline depletion by neonatal DSP-4: functional consequences and effect on a coexisting neurotransmitter (181) 147
Dornand, J., see El-Moatassim, C. (181) 111
Dray, A., Bettaney, J., Rueff, A., Walpole, C. and Wrigglesworth, R.
NE-19550 and NE-21610, antinociceptive capsaicin analogues: studies on nociceptive fibres of the neonatal rat tail in vitro (181) 289

El-Fakahany, E.E., see Fryer, A.D. (181) 51
El-Moatassim, C., Mani, J.-C. and Dornand, J.
Extracellular ATP^{4-} permeabilizes thymocytes not only to cations but also to low-molecular-weight solutes (181) 111
Eldar, M., see Cohen, A. (181) 311
Ellison, G., see See, R.E. (181) 175
Escorihuela, R.M., see Fernández-Teruel, A. (181) 35

Fernández-Teruel, A., Escorihuela, R.M., Boix, F. and Tobeña, A.
Picrotoxin changes the effects of imipramine and desipramine in rats in the forced swimming test (181) 35
Frey, H.-H., see Wahle, H. (181) 1
Fryer, A.D., El-Fakahany, E.E. and Jacoby, D.B.
Parainfluenza virus type 1 reduces the affinity of agonists for muscarinic receptors in guinea-pig lung and heart (181) 51

Gaggi, R. and Gianni, A.M.
Effects of calcium antagonists on biogenic amines in discrete brain areas (181) 187
Garty, N.B., see Leiba, H. (181) 71
Giachetti, A., see Maggi, C.A. (181) 105
Gianni, A.M., see Gaggi, R. (181) 187
Gibb, W.R.G., see Rose, S. (181) 97
Giuliani, S., see Maggi, C.A. (181) 105
Goodman, S.A., see Gould, A.B. (181) 225
Gould, A.B. and Goodman, S.A.
Effect of an angiotensin-converting enzyme inhibitor on blood pressure and erythropoiesis in rats (181) 225

Hadjiconstantinou, M. and Neff, N.H.
Differential recovery of dopamine synthetic enzymes following MPTP and the consequences of GM1 ganglioside treatment (181) 137

0014-2999/90/$03.50 © 1990 Elsevier Science Publishers B.V. (Biomedical Division)

Harron, D.W.G., see Spiers, J.P. (181) 235
Heijna, M.H., Padt, M., Hogenboom, F., Portoghese, P.S., Mulder, A.H. and Schoffelmeer, A.N.M.
Opioid receptor-mediated inhibition of dopamine and acetylcholine release from slices of rat nucleus accumbens, olfactory tubercle and frontal cortex (181) 267
Henderson, G., see Kennedy, C. (181) 315
Henry, J.-P., see Vasse, M. (181) 23
Hertting, G., see Allgaier, C. (181) 279
Herzog, A.G., see Kühn-Velten, W.N. (181) 151
Hirai, Y., see Katsuki, H. (181) 323
Hogenboom, F., see Heijna, M.H. (181) 267
Hoiting, B., Molleman, A., Nelemans, A. and Den Hertog, A.
P_2-purinoceptor-activated membrane currents and inositol tetrakisphosphate formation are blocked by suramin (181) 127
Horisaka, K., see Yamada, J. (181) 319
Horn, A.S., see Timmerman, W. (181) 253

Itoh, H., Lederis, K.P. and Rorstad, O.P.
Relaxation of isolated bovine coronary arteries by vasoactive intestinal peptide (181) 199
Iwata, H., see Matsuda, T. (181) 295

Jackson, C.B., see Sancilio, L.F. (181) 303
Jackson, E.A., see Rose, S. (181) 97
Jacoby, D.B., see Fryer, A.D. (181) 51
Janssen, P.A.J., see Xhonneux, R. (181) 261
Jarvis, G.E., see Kennedy, C. (181) 315
Jenner, P., see Rose, S. (181) 97

Kanda, T., see Matsuda, T. (181) 295
Kannisto, P., see Nilsson, C. (181) 247
Kanti Poddar, M., see Kumar Ray, S. (181) 159
Katsuki, H., Nakai, S., Hirai, Y., Akaji, K.-i., Kiso, Y. and Satoh, M.
Interleukin-1β inhibits long-term potentiation in the CA3 region of mouse hippocampal slices (181) 323
Kennedy, C., Jarvis, G.E. and Henderson, G.
The ligand rimcazole antagonises (+)SKF 10,047, but not (+)3-PPP, in the mouse isolated vas deferens (181) 315
Kim, H.S., see Altar, C.A. (181) 17
Kimura, I., see Yamada, J. (181) 319
Kiso, Y., see Katsuki, H. (181) 323
Korte, S.M. and Bohus, B.
The effect of ipsapirone on behavioural and cardiac responses in the shock-probe/defensive burying test in male rats (181) 307
Kühn-Velten, W.N., Herzog, A.G. and Müller, M.R.
Acute effects of anticonvulsant drugs on gonadotropin-stimulated and precursor-supported androgen production in the rat testis (181) 151
Kumar Ray, S. and Kanti Poddar, M.
Interaction of central serotonin and dopamine in the regulation of carbaryl-iduced tremor (181) 159

Lederis, K.P., see Itoh, H. (181) 199
Leiba, H., Garty, N.B., Schmidt-Sole, J., Piterman, O., Azrad, A. and Salomon, Y.
The melanocortin receptor in the rat lacrimal gland: a model system for the study of MSH (melanocyte stimulating hormone) as a potential neurotransmitter (181) 71
Lembeck, F., see Donnerer, J. (181) 147
Lindvall-Axelsson, M., see Nilsson, C. (181) 247

Maggi, C.A., Giuliani, S., Giachetti, A. and Meli, A.
The effect of MK-801 on the micturition reflex in anesthetized rats (181) 105
Mani, J.-C., see El-Moatassim, C. (181) 111
Marksteiner, J., Prommegger, R. and Sperk, G.
Effect of anticonvulsant treatment on kainic acid-induced increases in peptide levels (181) 241
Marsden, C.D., see Rose, S. (181) 97
Martínez, E., see Artigas, F. (181) 9
Matsuda, T., Kanda, T., Seong, Y.H., Baba, A. and Iwata, H.
p-Chlorophenylalanine attenuates the pituitary-adrenocortical response to 5-HT_{1A} receptor agonists in mice (181) 295
McEwen, B.S., see Mendelson, S.D. (181) 329
Meli, A., see Maggi, C.A. (181) 105
Mendelson, S.D. and McEwen, B.S.
Testosterone increases the concentration of [^{3}H]8-hydroxy-2-(di-n-propylamino)tetralin. Binding at 5-HT_{1A} receptors in the medial preoptic nucleus of the castrated male rat (181) 329
Mir, A.K., see Bouhelal, R. (181) 89
Molleman, A., see Hoiting, B. (181) 127
Mulder, A.H., see Heijna, M.H. (181) 267
Müller, M.R., see Kühn-Velten, W.N. (181) 151
Munson, Jr., H.R., see Sancilio, L.F. (181) 303

Nakai, S., see Katsuki, H. (181) 323
Namiki, M., see Uehara, A. (181) 141
Neff, N.H., see Hadjiconstantinou, M. (181) 137
Nelemans, A., see Hoiting, B. (181) 127
Nguyen, H., see Abdelrahman, A. (181) 83
Ni, H.
The pulmonary vascular responses to peripheral and central applications of acetylcholine (181) 167
Nicoletti, F., see Copani, A. (181) 327
Nilsson, C., Kannisto, P., Lindvall-Axelsson, M., Owman, C. and Rosengren, E.
The neuropeptides vasoactive intestinal polypeptide, peptide histidine isoleucine and neuropeptide Y modulate [^{3}H]noradrenaline release from sympathetic nerves in the choroid plexus (181) 247
Nomoto, M., see Rose, S. (181) 97

Okamiya, H., see Yamamoto, M. (181) 207
Okamura, K., see Uehara, A. (181) 141
Okumura, T., see Uehara, A. (181) 141
Owman, C., see Nilsson, C. (181) 247

Padt, M., see Heijna, M.H. (181) 267
Palacios, J.M., see Waeber, C. (181) 283
Pang, C.C.Y., see Abdelrahman, A. (181) 83
Pinkus, L.M., see Sancilio, L.F. (181) 303
Pinkus, L.M., see Waeber, C. (181) 283
Piterman, O., see Leiba, H. (181) 71
Portoghese, P.S., see Heijna, M.H. (181) 267
Prommegger, R., see Marksteiner, J. (181) 241
Protais, P., see Vasse, M. (181) 23

Rehavi, M., see Cohen, A. (181) 311
Reneman, R.S., see Xhonneux, R. (181) 261
Rorstad, O.P., see Itoh, H. (181) 199
Rose, S., Nomoto, M., Jackson, E.A., Gibb, W.R.G., Jenner, P. and Marsden, C.D.
1-Methyl-4-(2′-methylphenyl)-1,2,3,6-tetrahydropyridine (2′-methyl-MPTP) is less neurotoxic than MPTP in the common marmoset (181) 97
Rosengren, E., see Nilsson, C. (181) 247
Rueff, A., see Dray, A. (181) 289

Salomon, Y., see Leiba, H. (181) 71
Sancilio, L.F., Pinkus, L.M., Jackson, C.B. and Munson, Jr., H.R.
Emetic activity of zacopride in ferrets and its antagonism by pharmacological agents (181) 303
Satoh, M., see Katsuki, H. (181) 323
Schmidt, C.J. and Taylor, V.L.
Reversal of the acute effects of 3,4-methylenedioxymethamphetamine by 5-HT uptake inhibitors (181) 133
Schmidt, C.J., Black, C.K. and Taylor, V.L.
Antagonism of the neurotoxicity due to a single administration of methylenedioxymethamphetamine (181) 59
Schmidt-Sole, J., see Leiba, H. (181) 71
Schoffelmeer, A.N.M., see Heijna, M.H. (181) 267
Schuurkes, J.A.J., see Van den Brink, H.W. (181) 119
See, R.E. and Ellison, G.
Comparison of chronic administration of haloperidol and the atypical neuroleptics, clozapine and raclopride, in an animal model of tardive dyskinesia (181) 175
Segal, M.
Potassium currents activated in hippocampal neurons by serotonin are mediated by a change in intracellular calcium concentrations (181) 299
Seong, Y.H., see Matsuda, T. (181) 295
Shimizu, M., see Yamamoto, M. (181) 207
Skofitsch, G., see Donnerer, J. (181) 147
Sperk, G., see Marksteiner, J. (181) 241
Spiers, J.P., Harron, D.W.G. and Wilson, R.
Acute and chronic effects of rilmenidine on baroreflex function in conscious dogs (181) 235
Sugimoto, Y., see Yamada, J. (181) 319

Takasugi, Y., see Uehara, A. (181) 141
Takeuchi, N., see Yamada, J. (181) 319
Taylor, V.L., see Schmidt, C.J. (181) 59
Taylor, V.L., see Schmidt, C.J. (181) 133
Tepper, P.G., see Timmerman, W. (181) 253
Timmerman, W., Tepper, P.G., Bohus, B.G.J. and Horn, A.S.
The potential antipsychotic activity of the partial dopamine receptor agonist (+)N-0437 (181) 253
Tobeña, A., see Fernández-Teruel, A. (181) 35

Uehara, A., Okumura, T., Okamura, K., Takasugi, Y. and Namiki, M.
Lipopolysaccharide-induced inhibition of gastric acid and pepsin secretion in rats (181) 141

Van den Brink, H.W., Schuurkes, J.A.J., Van Nueten, J.M. and Van Rossum, J.M.
R 50 595, a selective non-competitive antagonist of cisapride, BRL 24924 and 5-hydroxytryptamine on the guinea-pig ileum (181) 119
Van Nueten, J.M., see Van den Brink, H.W. (181) 119
Van Rossum, J.M., see Van den Brink, H.W. (181) 119
Vasse, M., Chagraoui, A., Henry, J.-P. and Protais, P.
The rise of body temperature induced by the stimulation of dopamine D_1 receptors is increased in acutely reserpinized mice (181) 23

Waeber, C., Pinkus, L.M. and Palacios, J.M.
The (S)-isomer of [^{3}H]zacopride labels 5-HT_3 receptors with high affinity in rat brain (181) 283
Wahle, H. and Frey, H.-H.
Development of tolerance to the anticonvulsant effect of valproate but not to ethosuximide in a rat model of absence epilepsy (181) 1
Walpole, C., see Dray, A. (181) 289
Weizman, A., see Cohen, A. (181) 311
Wible, Jr., J.H. and DiMicco, J.A.
Resolution of central sites involved in picrotoxin-induced vagal activation and vasopressin release (181) 215
Wilson, R., see Spiers, J.P. (181) 235
Wouters, L., see Xhonneux, R. (181) 261
Wrigglesworth, R., see Dray, A. (181) 289

Xhonneux, R., Wouters, L., Reneman, R.S. and Janssen, P.A.J.
The l-enantiomer of nebivolol potentiates the blood pressure lowering effect of the d-enantiomer (181) 261

Yamada, J., Sugimoto, Y., Kimura, I., Takeuchi, N. and Horisaka, K.
The activation of serotonin receptors by tryptamine induces hyperinsulinemia in mice (181) 319
Yamamoto, M., Shimizu, M. and Okamiya, H.
Pharmacological actions of a new TRH analogue, YM-14673, in rats subjected to cerebral ischemia and anoxia (181) 207

Zafirova, M., see Carey, H.V. (181) 43